

## 3. Linear Algebra Review

- Range and Null space
- Left and right invertibility
- Rank
- Conservation of dimension
- Invertibility

## The Range

The range is defined to be

$$\text{range}(A) = \{ Ax \mid x \in \mathbb{R}^n \}$$

for *control* problems:

$$\begin{aligned} \text{range}(A) &= \text{set of possible outputs of } y = Ax \\ &= \text{span of columns of } A \end{aligned}$$

the range is also called the *column space* or the *image* of  $A$

the range is important in control problems because

The equation $y = Ax$ has a solution $x$ $\iff$ $y \in \text{range}(A)$
---


If we want a solution for all  $y \in \mathbb{R}^m$ , then we need  $\text{range}(A) = \mathbb{R}^m$ .

We call such a matrix *full-range* or *fully controllable* or *onto* or *surjective* or *right-invertible*

## Full Range Space

range  $A = \mathbb{R}^m$  is immediately equivalent to

- the columns of  $A$  span  $\mathbb{R}^m$ .
- there is a *right inverse function*  $h : \mathbb{R}^n \rightarrow \mathbb{R}^m$  so that  $Ah(y) = y$



Notice that  $h$  is a *controller* that gives an input that exactly produces  $y_{\text{des}}$ , since

$$y = Ax_{\text{applied}} = Ah(y_{\text{des}}) = y_{\text{des}}$$

We'll see that in fact there is a *linear* right inverse  $B$  such that  $AB = I$ .

Not so immediately, we'll see that  $\text{range } A = \mathbb{R}^m$  if and only if

- the rows of  $A$  are linearly independent.
- $AA^T$  is invertible.

## The Null Space

$$\text{null}(A) = \{ x \in \mathbb{R}^n \mid Ax = 0 \}$$

for *estimation* problems:

$$\begin{aligned} \text{null}(A) &= \text{set of unknowns which produce zero sensor output} \\ &= \text{set of vectors orthogonal to all rows of } A \end{aligned}$$

the null space is also called the *kernel* of  $A$ .

if  $z \in \text{null } A$  then  $x$  and  $x + z$  are indistinguishable from sensor reading  $y$

the null space is important in estimation problems because

if  $x_0$  is one solution to  $y = Ax$ , then the set of all solutions is

$$\{ x_0 + z \mid z \in \text{null}(A) \}$$


This follows immediately from the definition of null space, and means that we can *unambiguously* determine  $x$  from  $y$  if and only if  $\text{null}(A) = \{0\}$ .

We call  $A$  *zero-nullspace* or *fully observable* or *one-to-one* or *injective* or *left-invertible*

## Zero Null Space

$\text{null}(A) = \{0\}$  is immediately equivalent to

- the columns of  $A$  are linearly independent.
- there is a *left-inverse function*, a function  $g : \mathbb{R}^m \rightarrow \mathbb{R}^n$  so that  $g(Ax) = x$



Notice that the function  $g$  is an estimator that can exactly reconstruct  $x$ , because

$$x_{\text{est}} = g(y_{\text{meas}}) = g(Ax) = x$$

We'll see that there is a *linear* left inverse  $C$  such that  $CA = I$ .

We'll see that  $\text{null } A = \{0\}$  if and only if

- the rows of  $A$  span  $\mathbb{R}^n$ , that is,  $\text{range}(A^T) = \mathbb{R}^n$ .
- $A^T A$  is invertible

## Subspaces

A subset  $S \subset \mathbb{R}^n$  is called a *subspace* if

$$\begin{aligned} x + y &\in S \\ \lambda x &\in S \end{aligned}$$

for all  $x, y \in S$  and all  $\lambda \in \mathbb{R}$ .

- A subspace is closed under addition and scalar multiplication
- If  $A \in \mathbb{R}^{m \times n}$  then
  - $\text{range } A$  is a subspace of  $\mathbb{R}^m$
  - $\text{null } A$  is a subspace of  $\mathbb{R}^n$

These are *numerical representations* of a subspace

- Easy to test if  $x \in S$  with these representations. . .  
How to measure distance of  $x$  from  $S$ ?

## Rank

the *rank* of a matrix is the dimension of its column space. i.e.,

$$\text{rank}(A) = \dim \text{range}(A)$$

### Terminology

- an  $m \times n$  matrix  $A$  is called *full rank* if

$$\text{rank}(A) = \min\{m, n\}$$

- $A$  is called *skinny* if  $n < m$  and *fat* if  $n > m$
- $A$  is called *full column rank* if its columns are linearly independent


### Properties:

- Later: the dim. of the column space equals the dimension of the row space. i.e.,

$$\text{rank}(A) = \text{rank}(A^T)$$

## Coding interpretation of rank

- easy to see that the rank of product satisfies  $\text{rank } BC \leq \min\{\text{rank } B, \text{rank } C\}$
- hence if  $A = BC$  with  $B \in \mathbb{R}^{m \times r}$ ,  $C \in \mathbb{R}^{r \times n}$ , then  $\text{rank } A \leq r$
- conversely, we'll see later that if  $\text{rank } A = r$  then  $A \in \mathbb{R}^{m \times n}$  can be factored as  $A = BC$  with  $B \in \mathbb{R}^{m \times r}$ ,  $C \in \mathbb{R}^{r \times n}$



- $\text{rank}(A) = r$  is minimum size of vector needed to faithfully reconstruct  $y$  from  $x$

## Application: fast matrix-vector multiplication

- need to compute matrix-vector product  $y = Ax$ ,  $A \in \mathbb{R}^{m \times n}$
- $A$  has known factorization  $A = BC$ ,  $B \in \mathbb{R}^{m \times r}$
- computing  $y = Ax$  directly:  $mn$  operations
- computing  $y = Ax$  as  $y = B(Cx)$  (compute  $z = Cx$  first, then  $y = Bz$ ):  $rn + mr = (m + n)r$  operations
- savings can be considerable if  $r \ll \min\{m, n\}$

## Conservation of dimension

an important property of rank is

$$\dim \text{range}(A) + \dim \text{null}(A) = n$$

We interpret this as *conservation of dimension*. Of  $n$  input dimensions, every one is either mapped to zero or mapped to the output.

Therefore

- if  $\text{range } A = \mathbb{R}^m$  then  $n \geq m$ .
- if  $\text{null } A = \{0\}$  then  $n \leq m$ .
- and hence if both hold,  $n = m$ .

## Proof of conservation of dimension

Suppose  $A = [a_1 \ \dots \ a_n]$  where  $a_i \in \mathbb{R}^m$ . Then

$$\dim \text{range } A + \dim \text{null } A = n$$

*Proof:* Let  $V_1$  be a matrix whose columns are a basis for  $\text{null } A$ , and let  $V_2$  be a matrix whose columns complete that basis to one for  $\mathbb{R}^n$ . Let

$$V = [V_1 \ V_2]$$

so that  $\text{range } V = \mathbb{R}^n$ , then

$$\begin{aligned} \text{range } A &= \text{range } AV \\ &= \text{range } [0 \ AV_2] \\ &= \text{range } AV_2 \end{aligned}$$

since  $AV_1 = 0$ . Further, the columns of  $AV_2$  are linearly independent. To see this, suppose not, that is that there exists a nonzero vector  $z$  such that  $AV_2 z = 0$ . Then we would have  $V_2 z \in \text{null } A$ , which is impossible. Hence the columns of  $AV_2$  are a basis for  $\text{range } A$ , and there are  $n - \dim \text{null } A$  of them.

## Invertibility

If  $\text{range } A = \mathbb{R}^m$  and  $\text{null } A = \{0\}$  then

- $A$  is square, that is,  $m = n$
- $A$  has a left-inverse  $g(Ax) = x$  and a right-inverse  $Ah(y) = y$

In fact

- The inverses are equal,  $h = g$
- and linear, so there is a matrix  $B$  such that

$$AB = BA = I$$

- and  $B$  is the unique such matrix

Such a matrix is called *invertible*

## Invertibility

Suppose  $A$  is left-invertible and right-invertible. Then there is a matrix  $B$  such that

$$AB = BA = I$$

*Proof:* Since  $g(Ax) = x$  and  $Ah(y) = y$ , we have

$$g \circ A \circ h = h \quad g \circ A \circ g = g$$

Hence  $g = h$ . Call this function  $f$ . To see that  $f$  is linear, suppose  $Ax_1 = y_1$  and  $Ax_2 = y_2$ . Then we know

$$A(\alpha x_1 + \beta x_2) = \alpha y_1 + \beta y_2$$

and so

$$\alpha x_1 + \beta x_2 = f(\alpha y_1 + \beta y_2)$$

and so

$$\alpha f(y_1) + \beta f(y_2) = f(\alpha y_1 + \beta y_2)$$

which means  $f$  is linear.